Rehabilitation Protocol: Anterior Cruciate Ligament (ACL) Reconstruction with Hamstrings Autograft



Name:	Date:
Diagnosis:	Date of Surgery:
 Phase I (Weeks 0-4) Weightbearing: As tolerated with crutches (may be modifine transplant or articular cartilage procedure is performed) Hinged Knee Brace: Locked in full extension for ambulation and sleeping Unlocked for ambulation and removed while sleeping Range of Motion – AAROM → AROM as tolerated Therapeutic Exercises Quad/Hamstring sets and heel slides Non-weightbearing stretch of the Gastroc/Soleus Straight-Leg Raise with brace in full extension until No Hamstring Stretching Until 4 Weeks Post-Op 	g (Weeks 0-1) ng (Weeks 1-4) quad strength prevents extension lag
 Phase II (Weeks 4-6) Weightbearing: As tolerated discontinue crutch use Hinged Knee Brace: Discontinue brace use when patient leavension lag Range of Motion - Maintain full knee extension - work on Therapeutic Exercises Closed chain extension exercises Hamstring Stretching, Toe Raises, Balance Exercises Progress to weightbearing stretch of the Gastroc/So Begin use of the stationary bicycle 	n progressive knee flexion s
Phase III (Weeks 6-16) • Weightbearing: Full weightbearing • Range of Motion – Full/Painless ROM • Therapeutic Exercises • Begin Hamstring strengthening • Advance closed chain strengthening exercises, prop • Begin use of the Stairmaster/Elliptical • Can Start Straight Ahead Running at 12 Weeks Phase IV (Months 4-6) • Continue with strengthening (quad/hamstring) and • Begin cutting exercises and sport-specific drills	•
 Maintenance program for strength and endurance Return to sports at 6 months Comments:	
Frequency: times per week Duration: we Signature:	eeks Dato: