## **Rehabilitation Protocol: Medial Patellofemoral Ligament Reconstruction**



name:	Date:
Diagnosis:	Date of Surgery:
Phase I (Weeks 0-2)	
Weightbearing: Toe-touch weightbearing with crute	ches
Hinged Knee Brace:	
<ul> <li>Locked in full extension for ambulation and sl</li> </ul>	leening
Range of Motion – PROM/AAROM	reching
• Therapeutic Exercises	
Quad/Hamstring sets	
o Heel slides/Prone hangs/Patellar mobilization	n
<ul> <li>Straight-Leg Raise with brace in full extension</li> </ul>	
Phase II (Weeks 2-6)	
<ul> <li>Weightbearing: As tolerated – wean from crutch use</li> </ul>	e
<ul> <li>Hinged Knee Brace: Unlocked</li> </ul>	
<ul> <li>Range of Motion – PROM/AAROM/AROM: Maintain flexion (goal of 90 degrees by week 6)</li> </ul>	full knee extension – work on progressive kne
<ul> <li>Therapeutic Exercises</li> </ul>	
<ul> <li>LIMIT WEIGHBEARING EXERCISE TO FLEXIO</li> </ul>	N ANGLES < 90 DEGREES
<ul> <li>Isometric quadriceps strengthening</li> </ul>	
o Heel slides/Prone hangs	
<ul> <li>Patellar mobilization</li> </ul>	
<ul> <li>Phase III (Weeks 6-12)</li> <li>Weightbearing: Full weightbearing without crutche</li> <li>Discontinue Knee Brace – Convert to Patellar State</li> <li>Range of Motion – Advance to Full/Painless ROM</li> <li>Therapeutic Exercises         <ul> <li>Wall sits/Lunges</li> <li>Proprioception training</li> <li>Stationary bicycle</li> </ul> </li> </ul>	
Phase IV (Months 3-4)	
Advance closed chain strengthening – leg press, leg co	urls
Plyometric and proprioception training	
Treadmill jogging/Elliptical	
Phase V (Months 4-6)	
Gradual return to athletic activity as tolerated	
Maintenance program for strength and endurance	
Comments:	
Frequency: times per week	weeks
Signature:	Date: