Rehabilitation Protocol: Arthroscopic Meniscectomy/Chondral Debridement



| Name: | Date: |
|---|--------------------------------------|
| Diagnosis: | Date of Surgery: |
| | |
| | |
| Phase I (Weeks 0-2) |) 04 401 |
| Weightbearing: As tolerated with crutches (for balance) | ce) x 24-48 hours – progress to WBAT |
| • Range of Motion – AAROM → AROM as tolerated | |
| Goal: Immediate full range of motion | |
| Therapeutic Exercises | |
| Quad and Hamstring sets | |
| Heel slides | |
| Co-contractions | |
| Isometric adduction and abduction exercises | |
| Straight-leg raisesPatellar mobilization | |
| Patellar mobilization | |
| Phase II (Weeks 2-4) | |
| Weightbearing: As tolerated | |
| • Range of Motion – maintain full ROM – gentle passive | stretching at end ranges |
| • Therapeutic Exercises | stretening at ena ranges |
| Quadriceps and Hamstring strengthening | |
| Lunges | |
| o Wall-sits | |
| Balance exercises – Core work | |
| | |
| Phase III (Weeks 4-6) | |
| Weightbearing: Full weightbearing | |
| Range of Motion – Full/Painless ROM | |
| • Therapeutic Exercises | |
| o Leg press | |
| Hamstring curls | |
| o Squats | |
| Plyometric exercises | |
| Endurance work | |
| Return to athletic activity as tolerated | |
| | |
| Comments: | |
| | |
| Frequency: times per week Duration: | _ weeks |
| | |
| Signature: | Date: |